

Original Russian text www.bionet.nsc.ru/vogis/

The molecular view of mechanical stress of brain cells, local translation, and neurodegenerative diseases

T.M. Khlebodarova^{1, 2}

¹ Institute of Cytology and Genetics of Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia

² Kurchatov Genomic Center of the Institute of Cytology and Genetics of Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia

✉ tamara@bionet.nsc.ru

Abstract. The assumption that chronic mechanical stress in brain cells stemming from intracranial hypertension, arterial hypertension, or mechanical injury is a risk factor for neurodegenerative diseases was put forward in the 1990s and has since been supported. However, the molecular mechanisms that underlie the way from cell exposure to mechanical stress to disturbances in synaptic plasticity followed by changes in behavior, cognition, and memory are still poorly understood. Here we review (1) the current knowledge of molecular mechanisms regulating local translation and the actin cytoskeleton state at an activated synapse, where they play a key role in the formation of various sorts of synaptic plasticity and long-term memory, and (2) possible pathways of mechanical stress intervention. The roles of the mTOR (mammalian target of rapamycin) signaling pathway; the RNA-binding FMRP protein; the CYFIP1 protein, interacting with FMRP; the family of small GTPases; and the WAVE regulatory complex in the regulation of translation initiation and actin cytoskeleton rearrangements in dendritic spines of the activated synapse are discussed. Evidence is provided that chronic mechanical stress may result in aberrant activation of mTOR signaling and the WAVE regulatory complex via the YAP/TAZ system, the key sensor of mechanical signals, and influence the associated pathways regulating the formation of F actin filaments and the dendritic spine structure. These consequences may be a risk factor for various neurological conditions, including autistic spectrum disorders and epileptic encephalopathy. In further consideration of the role of the local translation system in the development of neuropsychic and neurodegenerative diseases, an original hypothesis was put forward that one of the possible causes of synaptopathies is impaired proteome stability associated with mTOR hyperactivity and formation of complex dynamic modes of *de novo* protein synthesis in response to synapse-stimulating factors, including chronic mechanical stress.

Key words: synapse; YAP/TAZ mechanosensor; mTOR; FMRP-dependent translation; complex dynamics; F actin; WAVE regulatory complex; autism spectrum disorders; epileptic encephalopathy.

For citation: Khlebodarova T.M. The molecular view of mechanical stress of brain cells, local translation, and neurodegenerative diseases. *Vavilovskii Zhurnal Genetiki i Seleksii = Vavilov Journal of Genetics and Breeding*. 2021;25(1): 92-100. DOI 10.18699/VJ21.011

Механический стресс клеток мозга, локальная трансляция и нейродегенеративные заболевания: молекулярно-генетические аспекты

Т.М. Хлебодарова^{1, 2}

¹ Федеральный исследовательский центр Институт цитологии и генетики Сибирского отделения Российской академии наук, Новосибирск, Россия

² Курчатовский геномный центр Института цитологии и генетики Сибирского отделения Российской академии наук, Новосибирск, Россия

✉ tamara@bionet.nsc.ru

Аннотация. Идея о том, что хронический механический стресс, который испытывают клетки мозга при повышенном внутричерепном давлении, артериальной гипертензии или вследствие травмы, может быть одним из факторов риска в развитии нейродегенеративных заболеваний, появилась еще в 90-е годы прошлого столетия и поддерживается в настоящее время. Однако молекулярно-генетические механизмы реализации событий, ведущих от механического воздействия на клетки к нарушению пластичности синапсов и последующему изменению поведения, когнитивных способностей и памяти, не ясны. В настоящем обзоре рассмотрены существующие данные о молекулярно-генетических механизмах регуляции локальной трансляции и актинового цитоскелета в активированном синапсе, играющих центральную роль в формировании различных видов пластичности синапса и долговременной памяти, и возможных путях влияния механического стресса на их состояние. Обсуждается роль mTOR сигнального каскада, РНК-связывающего белка FMRP, белка CYFIP1, взаимодействующего с FMRP, семейства малых ГТФаз и WAVE регуляторного комплекса

в регуляции инициации локальной трансляции и перестроек актинового цитоскелета в дендритных шипиках активированного синапса. Приводятся факты, свидетельствующие о том, что в условиях хронического механического стресса возможна aberrантная активация mTOR сигнального каскада и WAVE регуляторного комплекса через сенсор механических сигналов – регуляторный фактор YAP/TAZ, следствием которой могут быть нарушения активности системы локальной трансляции, а также связанных с ними механизмов регуляции формирования F-активных филаментов и структуры дендритных шипиков. Это может быть одной из причин развития различных неврологических патологий, включая аутистические расстройства и эпилептическую энцефалопатию. Высказывается оригинальная гипотеза, согласно которой одной из возможных причин синаптопатий может быть нарушение стабильности протеома, связанное с гиперактивностью mTOR и формированием сложных динамических режимов синтеза белков *de novo* в ответ на стимуляцию синапса, в том числе и в условиях хронического механического стресса.

Ключевые слова: синапс; механосенсор YAP/TAZ; mTOR; FMRP-зависимая трансляция; сложная динамика; F-актин; WAVE регуляторный комплекс; расстройства аутистического спектра; эпилептическая энцефалопатия.

Mechanical stress and neurodegenerative disorders

Mechanical signals are an important factor that determines the fate of cells, including their proliferation, survival, and differentiation, and takes part in tissue regeneration and wound healing. Mechanotransduction involves the reception of these forces and their conversion to biochemical and molecular signals, in particular, triggering of signaling pathways and expression of certain genes to allow cell adaptation to physical environment. There is ample evidence for the central role of the transcription regulator YAP (yes-associated protein 1) and its paralog TAZ (transcriptional co-activator with PDZ-binding motif), collectively named YAP/TAZ, as mechanical signal sensors and mediators (Dupont et al., 2011; Totaro et al., 2018; Dasgupta, McCollum, 2019). Impaired interaction of a cell and its environment causes aberrant YAP/TAZ activation and eventually a variety of diseases: atherosclerosis, fibrosis, lung hypertension, inflammation, muscle dystrophy, and cancer (Levy Nogueira et al., 2015, 2018; Yu et al., 2015; Panciera et al., 2017; Hong et al., 2019; Zhu et al., 2020). Recent studies indicate that mechanical stress may be among the causes of neurodegenerative processes in the brain, e.g., Alzheimer's disease (Levy Nogueira et al., 2015, 2016a, b, 2018).

The assumption that chronic mechanical stress experienced by brain cells exposed to intracranial hypertension, arterial hypertension, or mechanical injury is a risk factor for Alzheimer's disease and other neurodegenerative conditions was put forward as early as (Wostyn, 1994), and it is supported still now (Levy Nogueira et al., 2018).

What facts point to the existence of mechanisms by which mechanical stress influences nerve cell functions? First, it has been found that YAP/TAZ, being the key sensor and mediator of mechanical signals, activates the mTOR (mammalian target of rapamycin) signaling pathway (Tumaneng et al., 2012; McCarthy, 2013; Hu et al., 2017). This pathway is the central regulator of cap-dependent translation at a synapse. This regulation supports the dynamic plasticity of the synapse in response to external stimuli, and this plasticity underlies learning and memory (Costa-Mattioli et al., 2009; Buffington et al., 2014; Rosenberg et al., 2014; Santini et al., 2014). Disruption of these processes causes synaptic dysfunction and various neuropsychic disorders (Trifonova et al., 2017). YAP/TAZ activates mTOR by two mechanisms, illustrated

in Fig. 1: by stimulating the transcription of Rheb GTPase (Ras homologue enriched in brain) (Hu et al., 2017), which is an mTORC1 kinase activator, and by inhibiting the translation of PTEN (phosphatase and tensin homolog) with miR29 microRNA, thereby inducing aberrant PI3K-mediated activation of mTORC1 and mTORC2 kinases (Tumaneng et al., 2012; McCarthy, 2013).

Second, the actin cytoskeleton is the key mediator of mechanical signals (Seo, Kim, 2018). Its rearrangements in dendritic spines contribute much to learning and long-term memory formation (Basu, Lamprecht, 2018; Borovac et al., 2018). They are controlled by Rho GTPases (Tapon, Hall, 1997), whose hypo- or hyperactivity results in dendritic spine structure distortion, defective memory, and poor learnability. It may also cause multiple neurodevelopment disorders of various origins (Ba et al., 2013; Pyronneau et al., 2017; Zamboni et al., 2018; Nishiyama, 2019). The function of Rho GTPases at an activated synapse depends considerably on their *de novo* synthesis, which is determined by mTOR activity (Briz et al., 2015).

The activation of the mTOR signaling pathway by the YAP/TAZ mechanosensor under mechanical stress (Tumaneng et al., 2012; McCarthy, 2013; Hu et al., 2017) also promotes the activation of the heteropentameric WAVE regulatory complex (WASP family verprolin homologue) via S6K kinase and RAC1 GTPase (Derivery et al., 2009) by inducing its breakdown into subcomplexes and interaction of WAVE1 with Arp2/3 (Cory, Ridley, 2002; Millard et al., 2004; Abekhoukh, Bardoni, 2014; Molinie, Gautreau, 2018). These processes result in aberrant actin polymerization and structural anomalies of dendritic spines (see Fig. 1).

Thus, the pathways of the influence of mechanical stimuli on nerve cell functioning may involve the activation of mTOR signaling and rearrangements of the actin cytoskeleton in dendritic spines, which, in turn, depend on the activity of the local translation system at the synapse, controlled by mTOR. Just the disturbances in the local translation system at the synapse, including those caused by enhanced mTOR activity (manifesting themselves as synapse plasticity aberrations in the form of imbalance between synapse excitation and inhibition (Gobert et al., 2020)) are thought to be associated with various neuropsychic conditions, including autism spectrum disorders (ASDs), epilepsy, Parkinson's disease,




Fig. 1. Possible pathways of the effect of mechanical stress mediated by mTOR signaling on the intensity of local translation and the formation of actin cytoskeleton in dendritic spines of glutamatergic synapses in pyramidal cells of the hippocampus.

mGluR – receptor protein; PIKE (PI3-kinase enhancer), Rheb (Ras homologue enriched in brain), and Rac1 – GTPases; PI3K – phosphatidylinositol-3-kinase; Akt – protein kinase B; TSC1/2 – tuberous sclerosis complex 1/2; mTOR (mechanistic target of rapamycin) – serine/threonine kinase; S6K1 – S6 kinase 1; PTEN – phosphatase and tensin homolog; PP2A – protein phosphatase 2A; YAP/TAZ – mechanosensor; WAVE-1 – WAVE-1 regulatory complex; Arp2/3 – actin binding proteins. Proteins whose gene mutations are associated with neurological disorders are shown in red. Green arrows indicate translation activation via PP2A phosphatase and actin polymerization via S6 kinase and Rac1 GTPase in response to synapse stimulation by glutamate. Red arrows indicate possible mechanisms by which mechanical stress affects mTOR signaling.

and Alzheimer's disease (Gkogkas, Sonenberg, 2013; Meng et al., 2013; Won et al., 2013; Cai et al., 2015; Huber et al., 2015; Prampano et al., 2015; Klein et al., 2016; Martin, 2016; Onore et al., 2017).

In this regard, the molecular mechanisms regulating local translation and dynamic rearrangements of the actin cytoskeleton in dendritic spines affected by this regulation enjoy close attention (Bramham, 2008).

Local translation and neurodegenerative disorders

There is convincing evidence that local cap-dependent translation in the postsynaptic space of a dendritic spine enables its dynamic plasticity in response to external stimuli, which underlies learning and memory (Huber et al., 2000; Costa-Mattioli et al., 2009; Rosenberg et al., 2014; Santini et al., 2014; Louros, Osterweil, 2016).

Numerous examples have been reported that impaired local translation control at a synapse brings forth various neuropsychic disorders, including ASDs, epilepsy, Parkinson's disease, etc. (Gkogkas, Sonenberg, 2013; Buffington et al., 2014; Klein et al., 2016; Martin, 2016; Trifonova et al., 2017). Figure 2 illustrates the main regulatory events mediating the activation of local protein synthesis in dendritic spines of glutamatergic synapses of hippocampal pyramidal cells in response to the stimulation of metabotropic glutamate receptors (mGluR) on the postsynaptic membrane of excitatory synapses by glutamate. The local translation activity is controlled by the mTOR




Fig. 2. Schematic presentation of local translation regulation in dendritic spines of glutamatergic synapses of hippocampal pyramidal cells in response to synapse stimulation.

mGluR – receptor protein; NLGNs, Shank, PSD95, HOMERI – postsynaptic proteins; PIKE (PI3-kinase enhancer) and Rheb (Ras homologue enriched in brain) – GTPases; PI3K – phosphatidylinositol-3-kinase; Akt – protein kinase B; S6K1 – S6 kinase 1; TSC1/2 – tuberous sclerosis complex 1/2; mTOR (mechanistic target of rapamycin) – serine/threonine kinase; FMRP (fragile X mental retardation protein) – RNA binding protein; PP2A – protein phosphatase 2A; H-RAS – GTPase; RAF1, MEK1/2, ERK1/2 and MNK1/2 – kinases; eIF4E – factor of translation initiation; 4EBP2 – 4E-binding protein; CYFIP1 – cytoplasmic FMRP interacting protein 1. The proteinaceous products of genes whose mutations are associated with neurological disorders are shown in red. Green arrows indicate pathways of local translation activation via PP2A phosphatase and the RAS/ERK signaling pathway. The red arrow indicates blocking via the mTOR signaling pathway.

and RAS/ERK pathways (Huber et al., 2000; Darnell, Klann, 2013; Beggs et al., 2015; Chen, Joseph, 2015).

The key element in the regulation of local cap-dependent translation at a synapse is the RNA-binding fragile X mental retardation protein, or FMRP (Feng et al., 1997). It is the target of S6 kinase and PP2A phosphatase, which are activated in response to the stimulation of mGluR receptors (Narayanan et al., 2007, 2008). When phosphorylated, it arrests translation by binding to mRNA, ribosomes, and the eIF4E translation factor (Brown et al., 1998; Napoli et al., 2008; Chen et al., 2014). Dephosphorylation disrupts FMRP linkage to its targets, resulting in, on the one hand, to accelerated mRNA translation and, on the other hand, rapid degradation of FMRP itself (Nalavadi et al., 2012). FMRP controls translation efficiency through RNA binding sites (Chen, Joseph, 2015). It directly binds to the coding and 3'-UTR mRNA sequences (Brown et al., 1998; Darnell et al., 2011) and to the L5 protein of 80S ribosomes (Chen et al., 2014). In this way, it controls transcription elongation and termination. Translation can also be repressed in 3'-UTR by physical interaction of FMRP with the 43-kDa TAR DNA-binding protein (TDP-43) (Majumder et al., 2016).

FMRP is also involved in translation regulation at its initiation step by interaction with the cytoplasmic FMRP-interacting protein 1 (CYFIP1) (Napoli et al., 2008). The current notions of mechanisms regulating translation by means of FMRP (Napoli et al., 2008; Majumder et al., 2016) presume the interaction of a single molecule of the protein with 3'-UTR via TDP-43 and with translation initiation factor eIF4E via CYFIP1. Thus, FMRP and CYFIP1 are the key regulators of translation regulation at an activated synapse.

FMRP targets are mRNAs for proteinaceous components of the mTOR signaling pathway (PI3K kinase, PTEN phosphatase, tuberous sclerosis complex 2 (TSC2 and mTOR), PP2A phosphatase, receptor proteins (mGluR, NMDAR, and AMPAR), proteins forming the postsynaptic membrane (NLGN, SHANK, and PSD95), the ubiquitin-dependent protein degradation system (E3 ubiquitin ligase), and its own mRNA (FMR1) (Brown et al., 1998; Muddashetty et al., 2007; Gross et al., 2010; Sharma et al., 2010; Darnell et al., 2011; Ascano et al., 2012). Apparently, FMRP plays the key role in dynamic proteome regulation at an activated synapse (Zukin et al., 2009; Iacoangeli, Tiedge, 2013).

It is known that mutations in genes encoding most of these proteins result in synapse malfunction and various disorders. Mutations in the gene for the SHANK3 protein of the postsynaptic membrane cause Phelan–McDermid syndrome; in the gene for PTEN phosphatase, Cowden's disease; for NF1, type 1 neurofibromatosis; in the genes for GTPase, H-RAS, RAF1, and MEK1 kinase, Costello and Noonan syndromes; TSC2-TSC1, tuberous sclerosis; FMRP, fragile X syndrome; UBE3A ubiquitin-protein ligase, Angelman syndrome; and in genes for neuroligins NLGN3/4 and neurexin NRXN1, typical autism (Trifonova et al., 2016). Mutations in the *Shank3* gene and its abnormal expression are also considered to cause autism, schizophrenia, and epilepsy (Peça et al., 2011; Mei et al., 2016; de Sena Cortabitarte et al., 2017; Monteiro, Feng, 2017; Fu et al., 2020). Mutations in the gene for PTEN phosphatase often bring forth various neurological diseases: macrocephaly, epilepsy, mental deficiency, and autism (Zhou, Parada, 2012; Trifonova et al., 2016).

These data suggest that synapse malfunctions are related to anomalies in local translation regulation. One of the possible synaptopathy causes is disturbed proteome stability, which hampers the formation of synapse plasticity and long-term memory (Cajigas et al., 2010). Indeed, just poor proteome stability is reported to be associated with autism and other neuropsychic disorders (Klein et al., 2016; Louros, Osterweil, 2016).

It should be mentioned that the structure-functional organization of the system regulating FMRP activity includes negative and positive feedback loops, which are instability factors in molecular systems (Mackey, Glass, 1977; Decroly, Goldbeter, 1982; Goldbeter et al., 2001; Bastos de Figueiredo et al., 2002; Likhoshvai et al., 2013, 2015, 2016, 2020; Kogai et al., 2015, 2017; Suzuki et al., 2016; Khlebodarova et al., 2017).

These regulatory loops act in different time spans. They are associated with rapid (ca. 1 min) translation activation of FMRP-dependent mRNAs via PP2A phosphatase and its rather

rapid (2–5 min) arrest via the activation of S6 kinase (Narayanan et al., 2007, 2008). That is, the normal work of a synapse is supported by fine dynamic interplay among components of these signaling pathways at an activated synapse (see Fig. 2).

Analysis of dynamic features of the local translation system shows that an increase in the rate and efficiency of FMRP-dependent translation may induce instability in the local translation system, in particular, just in the physiological range of its operation (Khlebodarova et al., 2018; Likhoshvai, Khlebodarova, 2019). This result suggests that the known cases of ASDs related to the hyperactivity of the translation system at synapses (Pramparo et al., 2015; Onore et al., 2017) stem from proteome stability impairments associated with the formation of complex dynamic patterns of receptor protein synthesis in response to synapse stimulation (Khlebodarova et al., 2018, 2020). It is a brand-new insight into possible causes of synaptopathies.

It should be added that the elevated activity of mTOR signaling is a feature of not only ASDs but also other psychic and neurological diseases: Alzheimer's disease (Pei, Hugon, 2008), epilepsy (Wong, 2010), and even Down syndrome (Troca-Marin et al., 2012). It is also presumed that elevated mTOR activity causes early senescence and age-related neurodegenerative conditions in humans (Johnson et al., 2013).

In this regard, the hypothesis that the high copy numbers of rRNAs in some individuals are a risk factor for the development of ASDs, schizophrenia, and mental deficiency appears to be reasonable (Chestkov et al., 2018; Porokhovnik, 2019; Porokhovnik, Lyapunova, 2019) on the assumption that individual variations in copy numbers of rRNA genes correlate with ribosome concentrations in a cell and the activity of the translational machinery.

The actin cytoskeleton and neurodegenerative diseases

The actin cytoskeleton structure determines the morphology of dendritic spines in nerve cells. Its rearrangements by rapid assembly of actin monomers (G actin) to filaments (F actin) and inverse disassembly are essential for the formation of synaptic plasticity and long-term memory (Penzes, Rafałovich, 2012; Basu, Lamprecht, 2018). Disturbances in the mechanisms regulating the formation of F actin filaments and dendritic spine structure are thought to be associated with neurodegenerative disorders: Alzheimer's disease, schizophrenia, and autism (Bamburg, Bernstein, 2016; Borovac et al., 2018; Forrest et al., 2018; Ben Zablah et al., 2020; Lauterborn et al., 2020). Fig. 3 illustrates the major regulatory events underlying actin cytoskeleton rearrangements in dendritic spines of glutamatergic synapses in hippocampal pyramidal cells, which are activated in response to the action of glutamate on metabotropic glutamate receptors mGluR and ionotropic glutamate receptors NMDAR (N-methyl-d-aspartate receptor) on the postsynaptic membrane of excitatory synapses.

The induction of actin filament formation and filament stabilization at an activated synapse depends substantially on the activity of cofilin and the WAVE regulatory complex, which is controlled by S6K, LIMK1, and PAK1 kinases via signaling pathways mediated by the RAS family of small




Fig. 3. Schematic presentation of the regulation of actin cytoskeleton formation in dendritic spines of glutamatergic synapses of hippocampal pyramidal cells in response to synapse stimulation.

mGluR, NMDAR – receptor proteins; H-RAS, RHOA, RAC1, CDC42 – RAS GTPases; CaMKII, MEK1/2, RAF1, ERK1/2, S6K1, PAK1, ROCK, LIMK1 – kinases; CYFIP1 – cytoplasmic FMRP interacting protein 1; WAVE-1 – WAVE-1 regulatory complex; Arp2/3 – actin binding proteins; Cofilin – actin depolymerization factor. The proteinaceous products of genes whose mutations are associated with neurological disorders are shown in red. Green arrows indicate the pathways of actin polymerization regulation via mTOR and S6K1, black – via CaMKII signaling.

GTPases: H-RAS, RhoA, Rac1, and Cdc42 (Tapon, Hall, 1997; Rex et al., 2009; Ip et al., 2011; Chen et al., 2017; Schaks et al., 2018). The operation of these signaling pathways at an activated synapse depends greatly on fast *de novo* synthesis of Rho GTPases (Briz et al., 2015). Arrest of protein synthesis in dendritic spines of hippocampal cells completely suppresses the stimulation of RhoA GTPase, cofilin phosphorylation, and actin polymerization (Briz et al., 2015). A mutation in the *Fmr1* gene, which encodes the FMRP protein, the key local transcription regulator, completely suppresses the physiological stimulation of GTPase Rac1 and its effector PAK1 kinase, disrupting the stabilization of actin filaments at hippocampal cell synapses (Chen et al., 2010).

Proceeding from the above, the activity of RAS GTPases controlling the formation and stabilization of actin filaments in dendritic spines depends directly on their *de novo* synthesis, i. e., on the activity of mTOR and FMRP-dependent local translation. It is conjectured that unstable local translation (Khlebodarova et al., 2018, 2020; Likhoshvai, Khlebodarova, 2019), also results in hypo- or hyperactivity of RAS, which, in turn, causes aberrations in the structure of dendritic spines and neurological disorders associated therewith (Ba et al., 2013; Pyronneau et al., 2017; Zamboni et al., 2018; Nishiyama, 2019).

Rac1 GTPase plays the key role in the regulation of the heteropentameric WAVE regulatory complex. The activity of this GTPase depends much on S6K and mTOR1 kinases. Normally, the WAVE complex is inactive, but its interaction

with Rac GTPase induces its dissociation into two subcomplexes: CYFIP1-containing and WAVE1-containing (Derivery et al., 2009). The latter interacts with the Arp2/3 (actin-related proteins) complex and induces actin polymerization, as in Fig. 3 (Cory, Ridley, 2002; Millard et al., 2004; Abekhoukh, Bardoni, 2014; Molinie, Gautreau, 2018).

The disintegration of the WAVE complex and aberrant WAVE1 activation cause epileptic encephalopathy (Nakashima et al., 2018; Zhang et al., 2019; Zweier et al., 2019; Schaks et al., 2020). This may happen in cases of abnormal stoichiometric control of WAVE component synthesis (Abekhoukh et al., 2017) or mutations disrupting the interaction between WAVE1 and CYFIP2 (Nakashima et al., 2018; Zhang et al., 2019; Zweier et al., 2019; Schaks et al., 2020).

It should be noted that CYFIP1, being one of the main components of the WAVE regulatory complex, is also involved in translation regulation at its initiation step by interaction with the RNA-binding FMRP protein (Napoli et al., 2011). Thus, the mechanisms regulating local translation and actin cytoskeleton rearrangements in neural dendritic spines are additionally interlinked via the CYFIP1 protein (De Rubeis et al., 2013).

Conclusions

Analysis of presently available data shows the mechanisms regulating the local translation system at synapses and dynamic rearrangements of the actin cytoskeleton in dendritic spines of nerve cells, which play the central role in the formation of various types of synapse plasticity and long-term memory, are closely linked to each other and to the activity of the YAP/TAZ mechanosensor. This sensor can indirectly, via mTOR and S6K kinase, affect both translation efficiency and the state of actin filaments in dendritic spines (Tapon, Hall, 1997; Tumaneng et al., 2012; McCarthy, 2013; Reddy et al., 2013; Briz et al., 2015; Hu et al., 2017; Seo, Kim, 2018).

It is well substantiated that mTOR hyperactivity and functional aberrations in practically every component of the local translation system and of the machinery controlling rearrangements of the actin cytoskeleton in dendritic spines can cause numerous neurodevelopment disorders of various origins (Pei, Hugon, 2008; Wong, 2010; Johnson et al., 2013; Pramparo et al., 2015; Onore et al., 2017; Pyronneau et al., 2017; Trifonova et al., 2017; Nakashima et al., 2018; Nishiyama, 2019; Zhang et al., 2019).

Theoretical analysis of the dynamic features of local translation system operation presented in (Khlebodarova et al., 2018, 2020; Likhoshvai, Khlebodarova, 2019) suggests that one of the possible mechanisms of neurological disorders arising under chronic mechanical stress is the abnormal hyperactivity of mTOR and local translation at the synapse. It induces the dynamic instability of *de novo* protein synthesis at the activated synapse.

Thus, it is obvious that chronic mechanical stress may be one of the risk factors for synaptopathies and neurodegenerative diseases because of mTOR hyperactivation, which disturbs proteome stability, much needed for proper synapse plasticity and long-term memory (Klein et al., 2016; Louros, Osterweil, 2016).

References

- Abekhoush S., Bardoni B. CYFIP family proteins between autism and intellectual disability: links with Fragile X syndrome. *Front. Cell. Neurosci.* 2014;8:81. DOI 10.3389/fncel.2014.00081.
- Abekhoush S., Sahin H.B., Grossi M., Zongaro S., Maurin T., Madrigal I., Kazue-Sugioka D., Raas-Rothschild A., Doulazmi M., Carrera P., Stachon A., Scherer S., Drula Do Nascimento M.R., Trembleau A., Arroyo I., Szatmari P., Smith I.M., Milà M., Smith A.C., Giangrande A., Caillé L., Bardoni B. New insights into the regulatory function of CYFIP1 in the context of WAVE- and FMRP-containing complexes. *Dis. Model. Mech.* 2017;10(4):463-474. DOI 10.1242/dmm.025809.
- Ascano M. Jr., Mukherjee N., Bandaru P., Miller J.B., Nusbaum J.D., Corcoran D.L., Langlois C., Munschauer M., Dewell S., Hafner M., Williams Z., Ohler U., Tuschl T. FMRP targets distinct mRNA sequence elements to regulate protein expression. *Nature*. 2012;492:382-386. DOI 10.1038/nature11737.
- Ba W., van der Raadt J., Nadif Kasri N. Rho GTPase signaling at the synapse: implications for intellectual disability. *Exp. Cell Res.* 2013; 319(15):2368-2374. DOI 10.1016/j.yexcr.2013.05.033.
- Bamburg J.R., Bernstein B.W. Actin dynamics and cofilin-actin rods in Alzheimer disease. *Cytoskeleton (Hoboken)*. 2016;73(9):477-497. DOI 10.1002/cm.21282.
- Bastos de Figueiredo J.C., Diambra L., Glass L., Malta C.P. Chaos in two-looped negative feedback systems. *Phys. Rev. E Stat. Nonlin. Soft. Matter Phys.* 2002;65:051905.
- Basu S., Lamprecht R. The role of actin cytoskeleton in dendritic spines in the maintenance of long-term memory. *Front. Mol. Neurosci.* 2018;11:143. DOI 10.3389/fnmol.2018.00143.
- Beggs J.E., Tian S., Jones G.G., Xie J., Iadevaia V., Jenei V., Thomas G., Proud C.G. The MAP kinase-interacting kinases regulate cell migration, vimentin expression and eIF4E/CYFIP1 binding. *Biochem J.* 2015;467(1):63-76. DOI 10.1042/BJ20141066.
- Ben Zablah Y., Merovitch N., Jia Z. The role of ADF/cofilin in synaptic physiology and Alzheimer's disease. *Front. Cell Dev. Biol.* 2020; 8:594998. DOI 10.3389/fcell.2020.594998.
- Borovac J., Bosch M., Okamoto K. Regulation of actin dynamics during structural plasticity of dendritic spines: Signaling messengers and actin-binding proteins. *Mol. Cell. Neurosci.* 2018;91:122-130. DOI 10.1016/j.mcn.2018.07.001.
- Bramham C.R. Local protein synthesis, actin dynamics, and LTP consolidation. *Curr. Opin. Neurobiol.* 2008;18(5):524-531. DOI 10.1016/j.conb.2008.09.013.
- Briz V., Zhu G., Wang Y., Liu Y., Avetisyan M., Bi X., Baudry M. Activity-dependent rapid local RhoA synthesis is required for hippocampal synaptic plasticity. *J. Neurosci.* 2015;35(5):2269-2282. DOI 10.1523/JNEUROSCI.2302-14.2015.
- Brown V., Small K., Lakkis L., Feng Y., Gunter C., Wilkinson K.D., Warren S.T. Purified recombinant Fmrp exhibits selective RNA binding as an intrinsic property of the fragile X mental retardation protein. *J. Biol. Chem.* 1998;273(25):15521-15527. DOI 10.1074/jbc.273.25.15521.
- Buffington S.A., Huang W., Costa-Mattioli M. Translational control in synaptic plasticity and cognitive dysfunction. *Annu. Rev. Neurosci.* 2014;37:17-38. DOI 10.1146/annurev-neuro-071013-014100.
- Cai Z., Chen G., He W., Xiao M., Yan L.J. Activation of mTOR: a culprit of Alzheimer's disease? *Neuropsychiatr. Dis. Treat.* 2015;11: 1015-1030. DOI 10.2147/NDT.S75717.
- Cajigas J.J., Will T., Schuman E.M. Protein homeostasis and synaptic plasticity. *EMBO J.* 2010;29:2746-2752. DOI 10.1038/emboj.2010.173.
- Chen B., Chou H.T., Brautigam C.A., Xing W., Yang S., Henry L., Doolittle L.K., Walz T., Rosen M.K. Rac1 GTPase activates the WAVE regulatory complex through two distinct binding sites. *Elife*. 2017;6:e29795. DOI 10.7554/elife.29795.
- Chen E., Joseph S. Fragile X mental retardation protein: A paradigm for translational control by RNA-binding proteins. *Biochimie*. 2015; 114:147-154. DOI 10.1016/j.biochi.2015.02.005.
- Chen E., Sharma M.R., Shi X., Agrawal R.K., Joseph S. Fragile X mental retardation protein regulates translation by binding directly to the ribosome. *Mol. Cell.* 2014;54:407-417. DOI 10.1016/j.molcel.2014.03.023.
- Chen L.Y., Rex C.S., Babayan A.H., Kramár E.A., Lynch G., Gall C.M., Lauterborn J.C. Physiological activation of synaptic Rac>PAK (p-21 activated kinase) signaling is defective in a mouse model of fragile X syndrome. *J. Neurosci.* 2010;30(33):10977-10984. DOI 10.1523/JNEUROSCI.1077-10.2010.
- Chestkov I.V., Jestkova E.M., Ershova E.S., Golimbet V.E., Lezheiko T.V., Kolesina N.Y., Porokhovnik L.N., Lyapunova N.A., Izhevskaya V.L., Kutsev S.I., Veiko N., Kostyuk S.V. Abundance of ribosomal RNA gene copies in the genomes of schizophrenia patients. *Schizophr. Res.* 2018;197:305-314. DOI 10.1016/j.schres.2018.01.001.
- Cory G.O.C., Ridley A.J. Cell motility: braking WAVEs. *Nature*. 2002; 418:732-733. DOI 10.1038/418732a.
- Costa-Mattioli M., Sossin W.S., Klann E., Sonnenberg N. Translational control of long-lasting synaptic plasticity and memory. *Neuron*. 2009;61:10-26. DOI 10.1016/j.neuron.2008.10.055.
- Darnell J.C., Klann E. The translation of translational control by FMRP: therapeutic targets for FXS. *Nat. Neurosci.* 2013;16(11): 1530-1536. DOI 10.1038/nn.3379.
- Darnell J.C., Van Driesche S.J., Zhang C., Hung K.Y., Mele A., Fraser C.E., Stone E.F., Chen C., Fak J.J., Chi S.W., Licatalosi D.D., Richter J.D., Darnell R.B. FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell*. 2011;146: 247-261. DOI 10.1016/j.cell.2011.06.013.
- Dasgupta I., McCollum D. Control of cellular responses to mechanical cues through YAP/TAZ regulation. *J. Biol. Chem.* 2019;294(46): 17693-17706. DOI 10.1074/jbc.REV119.007963.
- De Rubeis S., Pasciuto E., Li K.W., Fernández E., Di Marino D., Buzzi A., Ostroff L.E., Klann E., Zwartkruis F.J., Komiyama N.H., Grant S.G., Poujol C., Choquet D., Achsel T., Posthuma D., Smit A.B., Bagni C. CYFIP1 coordinates mRNA translation and cytoskeleton remodeling to ensure proper dendritic spine formation. *Neuron*. 2013;79(6):1169-1182. DOI 10.1016/j.neuron.2013.06.039.
- de Sena Cortabitarte A., Degenhardt F., Strohmaier J., Lang M., Weiss B., Roeth R., Giegling I., Heilmann-Heimbach S., Hofmann A., Rujescu D., Fischer C., Rietschel M., Nöthen M.M., Rapold G.A., Berkel S. Investigation of SHANK3 in schizophrenia. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 2017;174(4):390-398. DOI 10.1002/ajmg.b.32528.
- Decroly O., Goldbeter A. Bhythmicity, chaos, and other patterns of temporal self-organization in a multiply regulated biochemical system. *Proc. Natl. Acad. Sci. USA*. 1982;79:6917-6921. DOI 10.1073/pnas.79.22.6917.
- Derivery E., Lombard B., Loew D., Gautreau A. The Wave complex is intrinsically inactive. *Cell Motil. Cytoskeleton*. 2009;66(10):777-790. DOI 10.1002/cm.20342.
- Dupont S., Morsut L., Aragona M., Enzo E., Giulitti S., Cordenonsi M., Zanconato F., Le Digabel J., Forcato M., Bicciato S., Elvassore N., Piccolo S. Role of YAP/TAZ in mechanotransduction. *Nature*. 2011; 474(7350):179-183. DOI 10.1038/nature10137.
- Feng Y., Absher D., Eberhart D.E., Brown V., Malter H.E., Warren S.T. FMRP associates with polyribosomes as an mRNP, and the I304N mutation of severe fragile X syndrome abolishes this association. *Mol. Cell*. 1997;1(1):109-118. DOI 10.1016/s1097-2765(00)80012-x.
- Forrest M.P., Parnell E., Penzes P. Dendritic structural plasticity and neuropsychiatric disease. *Nat. Rev. Neurosci.* 2018;19(4):215-234. DOI 10.1038/nrn.2018.16.

- Fu Y.J., Liu D., Guo J.L., Long H.Y., Xiao W.B., Xiao W., Feng L., Luo Z.H., Xiao B. Dynamic change of shanks gene mRNA expression and DNA methylation in epileptic rat model and human patients. *Mol. Neurobiol.* 2020;57(9):3712-3726. DOI 10.1007/s12035-020-01968-5.
- Gkogkas C.G., Sonenberg N. Translational control and autism-like behaviors. *Cell. Logist.* 2013;3:e24551. DOI 10.4161/cl.24551.
- Govert D., Schohl A., Kutsarova E., Ruthazer E.S. TORC1 selectively regulates synaptic maturation and input convergence in the developing visual system. *Dev. Neurobiol.* 2020. DOI 10.1002/dneu.22782.
- Goldbeter A., Gonze D., Houart G., Leloup J.C., Halloy J., Dupont G. From simple to complex oscillatory behavior in metabolic and genetic control networks. *Chaos.* 2001;11:247-260. DOI 10.1063/1.1345727.
- Gross C., Nakamoto M., Yao X., Chan C.B., Yim S.Y., Ye K., Warren S.T., Bassell G.J. Excess phosphoinositide 3-kinase subunit synthesis and activity as a novel therapeutic target in fragile X syndrome. *J. Neurosci.* 2010;30:10624-10638. DOI 10.1523/JNEUROSCI.0402-10.2010.
- Hong L., Li Y., Liu Q., Chen Q., Chen L., Zhou D. The Hippo signaling pathway in regenerative medicine. *Methods Mol. Biol.* 2019; 1893:353-370. DOI 10.1007/978-1-4939-8910-2_26.
- Hu J.K., Du W., Shelton S.J., Oldham M.C., DiPersio C.M., Klein O.D. An FAK-YAP-mTOR signaling axis regulates stem cell-based tissue renewal in mice. *Cell Stem Cell.* 2017;21(1):91-106. DOI 10.1016/j.stem.2017.03.023.
- Huber K.M., Kayser M.S., Bear M.F. Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science.* 2000;288(5469):1254-1257. DOI 10.1126/science.288.5469.1254.
- Huber K.M., Klann E., Costa-Mattioli M., Zukin R.S. Dysregulation of mammalian target of rapamycin signaling in mouse models of autism. *J. Neurosci.* 2015;35(41):13836-13842. DOI 10.1523/JNEUROSCI.2656-15.2015.
- Iacoangeli A., Tiedge H. Translational control at the synapse: role of RNA regulators. *Trends Biochem. Sci.* 2013;38(1):47-55. DOI 10.1016/j.tibs.2012.11.001.
- Ip C.K., Cheung A.N., Ngan H.Y., Wong A.S. p70 S6 kinase in the control of actin cytoskeleton dynamics and directed migration of ovarian cancer cells. *Oncogene.* 2011;30(21):2420-2432. DOI 10.1038/onc.2010.615.
- Johnson S.C., Rabinovitch P.S., Kaeberlein M. mTOR is a key modulator of ageing and age-related disease. *Nature.* 2013;493(7432): 338-345. DOI 10.1038/nature11861.
- Khlebodarova T.M., Kogai V.V., Fadeev S.I., Likhoshvai V.A. Chaos and hyperchaos in simple gene network with negative feedback and time delays. *J. Bioinform. Comput. Biol.* 2017;15(2):1650042. DOI 10.1142/S0219720016500426.
- Khlebodarova T.M., Kogai V.V., Likhoshvai V.A. On the dynamical aspects of local translation at the activated synapse. *BMC Bioinformatics.* 2020;21(Suppl. 11):258. DOI 10.1186/s12859-020-03597-0.
- Khlebodarova T.M., Kogai V.V., Trifonova E.A., Likhoshvai V.A. Dynamic landscape of the local translation at activated synapses. *Mol. Psych.* 2018;23(1):107-114. DOI 10.1038/mp.2017.245.
- Khlebodarova T.M., Likhoshvai V.A., Kogai V.V. On the chaotic potential of local translation at activated synapses. In: Mathematical Biology and Bioinformatics. Puschnino: IMPB RAS, 2018;7:e68.1-e68.6. DOI 10.17537/icmmb18.6. (in Russian)
- Klein M.E., Monday H., Jordan B.A. Proteostasis and RNA binding proteins in synaptic plasticity and in the pathogenesis of neuropsychiatric disorders. *Neural Plast.* 2016;2016:3857934. DOI 10.1155/2016/3857934.
- Kogai V.V., Khlebodarova T.M., Fadeev S.I., Likhoshvai V.A. Complex dynamics in alternative mRNA splicing: mathematical model. *Vychislitelnye Technologii = Computational Technologies.* 2015; 20(1):38-52. (in Russian)
- Kogai V.V., Likhoshvai V.A., Fadeev S.I., Khlebodarova T.M. Multiple scenarios of transition to chaos in the alternative splicing model. *Int. J. Bifurcat. Chaos.* 2017;27(2):1730006. DOI 10.1142/S0218127417300063.
- Lauterborn J.C., Cox C.D., Chan S.W., Vanderklish P.W., Lynch G., Gall C.M. Synaptic actin stabilization protein loss in Down syndrome and Alzheimer disease. *Brain Pathol.* 2020;30(2):319-331. DOI 10.1111/bpa.12779.
- Levy Nogueira M., da Veiga Moreira J., Baroncio G.F., Dubois B., Steyaert J.M., Schwartz L. Mechanical stress as the common denominator between chronic inflammation, cancer, and Alzheimer's disease. *Front. Oncol.* 2015;5:197. DOI 10.3389/fonc.2015.00197.
- Levy Nogueira M., Epelbaum S., Steyaert J.M., Dubois B., Schwartz L. Mechanical stress models of Alzheimer's disease pathology. *Alzheimers Dement.* 2016;12(3):324-333. DOI 10.1016/j.jalz.2015.10.005.
- Levy Nogueira M., Hamraz M., Abolhassani M., Bigan E., Lafitte O., Steyaert J.M., Dubois B., Schwartz L. Mechanical stress increases brain amyloid β , tau, and α -synuclein concentrations in wild-type mice. *Alzheimers Dement.* 2018;14(4):444-453. DOI 10.1016/j.jalz.2017.11.003.
- Levy Nogueira M., Lafitte O., Steyaert J.M., Bakardjian H., Dubois B., Hampel H., Schwartz L. Mechanical stress related to brain atrophy in Alzheimer's disease. *Alzheimers Dement.* 2016;12(1):11-20. DOI 10.1016/j.jalz.2015.03.005.
- Likhoshvai V.A., Fadeev S.I., Kogai V.V., Khlebodarova T.M. On the chaos in gene networks. *J. Bioinform. Comput. Biol.* 2013;11(1): 1340009. DOI 10.1142/S021972001340009X.
- Likhoshvai V.A., Golubyatnikov V.P., Khlebodarova T.M. Limit cycles in models of circular gene networks regulated by negative feedbacks. *BMC Bioinformatics.* 2020;21(Suppl. 11):255. DOI 10.1186/s12859-020-03598-z.
- Likhoshvai V.A., Khlebodarova T.M. On stationary solutions of delay differential equations: a model of local translation in synapses. *Matematicheskaya Biologiya i Bioinformatika = Mathematical Biology and Bioinformatics.* 2019;14(2):554-569. DOI 10.17537/2019.14.554. (in Russian)
- Likhoshvai V.A., Kogai V.V., Fadeev S.I., Khlebodarova T.M. Alternative splicing can lead to chaos. *J. Bioinform. Comput. Biol.* 2015; 13(1):1540003. DOI 10.1142/S021972001540003X.
- Likhoshvai V.A., Kogai V.V., Fadeev S.I., Khlebodarova T.M. Chaos and hyperchaos in a model of ribosome autocatalytic synthesis. *Sci. Rep.* 2016;6:38870. DOI 10.1038/srep38870.
- Louros S.R., Osterweil E.K. Perturbed proteostasis in autism spectrum disorders. *J. Neurochem.* 2016;139(6):1081-1092. DOI 10.1111/jnc.13723.
- Mackey M.C., Glass L. Oscillation and chaos in physiological control systems. *Science.* 1977;197(4300):287-289. DOI 10.1126/science.267326.
- Majumder P., Chu J.F., Chatterjee B., Swamy K.B., Shen C.J. Co-regulation of mRNA translation by TDP-43 and fragile X syndrome protein FMRP. *Acta Neuropathol.* 2016;132(5):721-738. DOI 10.1007/s00401-016-1603-8.
- Martin I. Decoding Parkinson's disease pathogenesis: the role of deregulated mRNA translation. *J. Parkinsons Dis.* 2016;6(1):17-27. DOI 10.3233/JPD-150738.
- McCarthy N. Signalling: YAP, PTEN and miR-29 size each other up. *Nat. Rev. Cancer.* 2013;13(1):4-5. DOI 10.1038/nrc3422.
- Mei Y., Monteiro P., Zhou Y., Kim J.A., Gao X., Fu Z., Feng G. Adult restoration of Shank3 expression rescues selective autistic-like phenotypes. *Nature.* 2016;530(7591):481-484. DOI 10.1038/nature16971.
- Meng X.F., Yu J.T., Song J.H., Chi S., Tan L. Role of the mTOR signalling pathway in epilepsy. *J. Neurol. Sci.* 2013;332(1-2):4-15. DOI 10.1016/j.jns.2013.05.029.
- Millard T.H., Sharp S.J., Machesky L.M. Signalling to actin assembly via the WASP (Wiskott-Aldrich syndrome protein)-family proteins

- and the Arp2/3 complex. *Biochem. J.* 2004;380(Pt. 1):1-17. DOI 10.1042/BJ20040176.
- Molinie N., Gautreau A. The Arp2/3 regulatory system and its deregulation in cancer. *Physiol. Rev.* 2018;98(1):215-238. DOI 10.1152/physrev.00006.2017.
- Monteiro P., Feng G. SHANK proteins: roles at the synapse and in autism spectrum disorder. *Nat. Rev. Neurosci.* 2017;18(3):147-157. DOI 10.1038/nrn.2016.183.
- Muddashetty R.S., Kelić S., Gross C., Xu M., Bassell G.J. Dysregulated metabotropic glutamate receptor-dependent translation of AMPA receptor and postsynaptic density-95 mRNAs at synapses in a mouse model of fragile X syndrome. *J. Neurosci.* 2007;27(20):5338-5348. DOI 10.1523/JNEUROSCI.0937-07.2007.
- Nakashima M., Kato M., Aoto K., Shiina M., Belal H., Mukaida S., Kuromada S., Sato A., Zerem A., Lerman-Sagie T., Lev D., Leong H.Y., Tsurusaki Y., Mizuguchi T., Miyatake S., Miyake N., Ogata K., Saitsu H., Matsumoto N. De novo hotspot variants in CYFIP2 cause early-onset epileptic encephalopathy. *Ann. Neurol.* 2018;83(4):794-806. DOI 10.1002/ana.25208.
- Nalavadi V.C., Muddashetty R.S., Gross C., Bassell G.J. Dephosphorylation-induced ubiquitination and degradation of FMRP in dendrites: a role in immediate early mGluR-stimulated translation. *J. Neurosci.* 2012;32(8):2582-2587. DOI 10.1523/JNEUROSCI.5057-11.2012.
- Napoli I., Mercaldo V., Boyl P.P., Eleuteri B., Zalfa F., De Rubeis S., Di Marino D., Mohr E., Massimi M., Falconi M., Witke W., Costa-Mattioli M., Sonenberg N., Achsel T., Bagni C. The fragile X syndrome protein represses activity-dependent translation through CYFIP1, a new 4E-BP. *Cell.* 2008;134(6):1042-1054. DOI 10.1016/j.cell.2008.07.031.
- Narayanan U., Nalavadi V., Nakamoto M., Pallas D.C., Ceman S., Bassell G.J., Warren S.T. FMRP phosphorylation reveals an immediate-early signaling pathway triggered by group I mGluR and mediated by PP2A. *J. Neurosci.* 2007;27(52):14349-14357. DOI 10.1523/JNEUROSCI.2969-07.2007.
- Narayanan U., Nalavadi V., Nakamoto M., Thomas G., Ceman S., Bassell G.J., Warren S.T. S6K1 phosphorylates and regulates fragile X mental retardation protein (FMRP) with the neuronal protein synthesis-dependent mammalian target of rapamycin (mTOR) signaling cascade. *J. Biol. Chem.* 2008;283(27):18478-18482. DOI 10.1074/jbc.C800055200.
- Nishiyama J. Plasticity of dendritic spines: Molecular function and dysfunction in neurodevelopmental disorders. *Psychiat. Clin. Neurosci.* 2019;73(9):541-550. DOI 10.1111/pcn.12899.
- Onore C., Yang H., Van de Water J., Ashwood P. Dynamic Akt/mTOR signaling in children with autism spectrum disorder. *Front. Pediatr.* 2017;5:43. DOI 10.3389/fped.2017.00043.
- Panciera T., Azzolini L., Cordenonsi M., Piccolo S. Mechanobiology of YAP and TAZ in physiology and disease. *Nat. Rev. Mol. Cell Biol.* 2017;18(12):758-770. DOI 10.1038/nrm.2017.87.
- Peça J., Feliciano C., Ting J.T., Wang W., Wells M.F., Venkataraman T.N., Lascola C.D., Fu Z., Feng G. Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature.* 2011;472(7344):437-442. DOI 10.1038/nature09965.
- Pei J.J., Hugon J. mTOR-dependent signalling in Alzheimer's disease. *J. Cell. Mol. Med.* 2008;12(6b):2525-2532. DOI 10.1111/j.1582-4934.2008.00509.x.
- Penzes P., Rafalovich I. Regulation of the actin cytoskeleton in dendritic spines. *Adv. Exp. Med. Biol.* 2012;970:81-95. DOI 10.1007/978-3-7091-0932-8_4.
- Porokhovnik L. Individual copy number of ribosomal genes as a factor of mental retardation and autism risk and severity. *Cells.* 2019;8(10):1151. DOI 10.3390/cells8101151.
- Porokhovnik L.N., Lyapunova N.A. Dosage effects of human ribosomal genes (rDNA) in health and disease. *Chromosome Res.* 2019;27(1-2):5-17. DOI 10.1007/s10577-018-9587-y.
- Pramparo T., Pierce K., Lombardo M.V., Carter Barnes C., Marinero S., Ahrens-Barbeau C., Murray S.S., Lopez L., Xu R., Courchesne E. Prediction of autism by translation and immune/inflammation co-expressed genes in toddlers from pediatric community practice. *JAMA Psychiatry.* 2015;72:386-394. DOI 10.1001/jamapsychiatry.2014.3008.
- Pyronneau A., He Q., Hwang J.Y., Porch M., Contractor A., Zukin R.S. Aberrant Rac1-cofilin signaling mediates defects in dendritic spines, synaptic function, and sensory perception in fragile X syndrome. *Sci. Signal.* 2017;10(504):eaan0852. DOI 10.1126/scisignal.aan0852.
- Reddy P., Deguchi M., Cheng Y., Hsueh A.J. Actin cytoskeleton regulates Hippo signaling. *PLoS One.* 2013;8(9):e73763. DOI 10.1371/journal.pone.0073763.
- Rex C.S., Chen L.Y., Sharma A., Liu J., Babayan A.H., Gall C.M., Lynch G. Different Rho GTPase-dependent signaling pathways initiate sequential steps in the consolidation of long-term potentiation. *J. Cell Biol.* 2009;186(1):85-97. DOI 10.1083/jcb.200901084.
- Rosenberg T., Gal-Ben-Ari S., Dieterich D.C., Kreutz M.R., Ziv N.E., Gundelfinger E.D., Rosenblum K. The roles of protein expression in synaptic plasticity and memory consolidation. *Front. Mol. Neurosci.* 2014;7:86. DOI 10.3389/fnmol.2014.00086.
- Santini E., Huynh T.N., Klann E. Mechanisms of translation control underlying long-lasting synaptic plasticity and the consolidation of long-term memory. *Prog. Mol. Biol. Transl. Sci.* 2014;122:131-167. DOI 10.1016/B978-0-12-420170-5.00005-2.
- Schaks M., Reinke M., Witke W., Rottner K. Molecular dissection of neurodevelopmental disorder-causing mutations in CYFIP2. *Cells.* 2020;9(6):1355. DOI 10.3390/cells9061355.
- Schaks M., Singh S.P., Kage F., Thomason P., Klünemann T., Steffen A., Blankenfeldt W., Stradal T.E., Insall R.H., Rottner K. Distinct interaction sites of RAC GTPase with WAVE regulatory complex have non-redundant functions *in vivo*. *Curr. Biol.* 2018;28(22):3674-3684.e6. DOI 10.1016/j.cub.2018.10.002.
- Seo J., Kim J. Regulation of Hippo signaling by actin remodeling. *BMB Rep.* 2018;51(3):151-156. DOI 10.5483/bmbrep.2018.51.3.012.
- Sharma A., Hoeffer C.A., Takayasu Y., Miyawaki T., McBride S.M., Klann E., Zukin R.S. Dysregulation of mTOR signaling in fragile X syndrome. *J. Neurosci.* 2010;30(2):694-702. DOI 10.1523/JNEUROSCI.3696-09.2010.
- Suzuki Y., Lu M., Ben-Jacob E., Onuchic J.N. Periodic, quasi-periodic and chaotic dynamics in simple gene elements with time delays. *Sci. Rep.* 2016;6:21037. DOI 10.1038/srep21037.
- Tapon N., Hall A. Rho, Rac and Cdc42 GTPases regulate the organization of the actin cytoskeleton. *Curr. Opin. Cell Biol.* 1997;9(1):86-92. DOI 10.1016/s0955-0674(97)80156-1.
- Totaro A., Panciera T., Piccolo S. YAP/TAZ upstream signals and downstream responses. *Nat. Cell Biol.* 2018;20(8):888-899. DOI 10.1038/s41556-018-0142-z.
- Trifonova E.A., Khlebodarova T.M., Grunenko N.E. Molecular mechanisms of autism as a form of synaptic dysfunction. *Vavilovskii Zhurnal Genetiki i Selektii = Vavilov Journal of Genetics and Breeding.* 2016;20(6):959-967. DOI 10.18699/VJ16.217. (in Russian)
- Trifonova E.A., Khlebodarova T.M., Grunenko N.E. Molecular mechanisms of autism as a form of synaptic dysfunction. *Russ. J. Genet.: Appl. Res.* 2017;7(8):869-877.
- Troca-Marin J.A., Alves-Sampaio A., Montesinos M.L. Deregulated mTOR-mediated translation in intellectual disability. *Prog. Neurobiol.* 2012;96(2):268-282. DOI 10.1016/j.pneurobio.2012.01.005.
- Tumaneng K., Schlegelmilch K., Russell R.C., Yimlaimai D., Basnett H., Mahadevan N., Fitamant J., Bardeesy N., Camargo F.D., Guan K.L. YAP mediates crosstalk between the Hippo and PI(3)K-TOR pathways by suppressing PTEN via miR-29. *Nat. Cell Biol.* 2012;14(12):1322-1329. DOI 10.1038/ncb2615.
- Won H., Mah W., Kim E. Autism spectrum disorder causes, mechanisms, and treatments: focus on neuronal synapses. *Front. Mol. Neurosci.* 2013;6:19. DOI 10.3389/fnmol.2013.00019.
- Wong M. Mammalian target of rapamycin (mTOR) inhibition as a potential antiepileptogenic therapy: From tuberous sclerosis to com-

- mon acquired epilepsies. *Epilepsia*. 2010;51(1):27-36. DOI 10.1111/j.1528-1167.2009.02341.x.
- Wostyn P. Intracranial pressure and Alzheimer's disease: a hypothesis. *Med. Hypotheses*. 1994;43(4):219-222. DOI 10.1016/0306-9877(94)90069-8.
- Yu F.X., Zhao B., Guan K.L. Hippo pathway in organ size control, tissue homeostasis, and cancer. *Cell*. 2015;163(4):811-828. DOI 10.1016/j.cell.2015.10.044.
- Zamboni V., Jones R., Umbach A., Ammoni A., Passafaro M., Hirsch E., Merlo G.R. Rho GTPases in intellectual disability: from genetics to therapeutic opportunities. *Int. J. Mol. Sci.* 2018;19:1821. DOI 10.3390/ijms19061821.
- Zhang Y., Lee Y., Han K. Neuronal function and dysfunction of CYFIP2: from actin dynamics to early infantile epileptic encephalopathy. *BMB Rep.* 2019;52(5):304-311. DOI 10.5483/BMBRep.2019.52.5.097.
- Zhou J., Parada L.F. PTEN signaling in autism spectrum disorders. *Curr. Opin. Neurobiol.* 2012;22(5):873-879. DOI 10.1016/j.conb.2012.05.004.
- Zhu T., Ma Z., Wang H., Jia X., Wu Y., Fu L., Li Z., Zhang C., Yu G. YAP/TAZ affects the development of pulmonary fibrosis by regulating multiple signaling pathways. *Mol. Cell. Biochem.* 2020; 475(1-2):137-149. DOI 10.1007/s11010-020-03866-9.
- Zukin R.S., Richter J.D., Bagni C. Signals, synapses, and synthesis: how new proteins control plasticity. *Front. Neural Circuits*. 2009; 3:14. DOI 10.3389/neuro.04.014.2009.
- Zweier M., Begemann A., McWalter K., Cho M.T., Abela L., Banika S., Behring B., Berger A., Brown C.W., Carneiro M., Chen J., Cooper G.M. Deciphering Developmental Disorders (DDD) Study, Finnila C.R., Guillen Sacoto M.J., Henderson A., Hüffmeier U., Jøset P., Kerr B., Lesca G., Leszinski G.S., McDermott J.H., Melter M.R., Monaghan K.G., Mostafavi R., Öunap K., Plecko B., Powis Z., Purcarin G., Reimand T., Riedhammer K.M., Schreiber J.M., Sirsi D., Wierenga K.J., Wojcik M.H., Papuc S.M., Steindl K., Sticht H., Rauch A. Spatially clustering *de novo* variants in CYFIP2, encoding the cytoplasmic FMRP interacting protein 2, cause intellectual disability and seizures. *Eur. J. Hum. Genet.* 2019;27(5):747-759. DOI 10.1038/s41431-018-0331-z.

Acknowledgements. This work was supported by the budget project No. 0259-2021-0009.

Conflict of interest. The author declares no conflict of interest.

Received October 19, 2020. Revised December 21, 2020. Accepted December 22, 2020.